



RECENT RESULTS FROM MINOS AND FUTURE PROSPECTS

Jonathan M. Paley for the MINOS Collaboration
Indiana University


Fermilab Users' Meeting
June 5, 2008

The MINOS Collaboration



Argonne - Arkansas Tech - Athens - Benedictine - Brookhaven - Caltech - Cambridge - Campinas - Fermilab - Harvard - IIT - Indiana - Minnesota-Twin Cities - Minnesota-Duluth - Oxford - Pittsburgh - Rutherford - São Paulo - South Carolina - Stanford - Sussex - Texam A&M - Texas-Austin - Tufts - UCL - Warsaw - William & Mary

Goals of the MINOS Experiment



- Make precise measurement of Δm^2 and $\sin^2(2\theta)$


$$P(\nu_\mu \rightarrow \nu_\mu) = 1 - \sin^2(2\theta) \sin^2(1.27 \Delta m^2 L/E)$$

- Confirm oscillations vs. other explanations (decay, decoherence)
- Secondary goals:
 - Search for subdominant $\nu_\mu \rightarrow \nu_e$
 - Search for sterile neutrinos
 - CPT tests
 - Atmospheric neutrino and cosmic ray studies

2007-08: Very Productive Year!


- 2 boxes opened (ν_μ CC and NC blind analyses)
- 6 theses
- Significant progress in understanding backgrounds and systematic uncertainties in all analyses

The MINOS Experiment




Identifying Events in MINOS

ν_μ CC event




Long μ track +
shower at vertex

ν_e CC event



Short event with
EM shower profile.

NC event





Short, diffuse event.

$$E_\nu = E_{\text{shower}} + E_{\mu,e}$$


$$\delta E_{\text{shower}} = 55\%/\sqrt{E} \quad \delta E_\mu = 6\% \text{ range, } 10\% \text{ curvature}$$

Producing Neutrinos at the Main Injector





- Mesons produced in $120\text{ GeV}/c$ $p + \text{graphite}$ target interactions are focused in two magnetic horns.
- ν beam energy is *tunable* by moving target position longitudinally w.r.t. the horn positions.
- In LE beam configuration, beam is composed of 92.9% ν_μ , 5.8% $\bar{\nu}_\mu$, and 1.3% ν_e and $\bar{\nu}_e$.

Predicting the Flux





- MINOS uses Fluka06 MC to predict the ν flux.
- Uncertainty on flux is $\sim 30\%$ due to lack of hadron production data.
- To improve our data-to-MC agreement, we tune the Fluka MC to ND energy spectra of different beam configurations.
- These beam-reweighted spectra are used in all analyses discussed today.

Measurement of Hadron Production off NuMI Target in MIPP





- Main Injector Particle Production (MIPP) is a fixed target experiment with beams of π , K and p from 5-120 GeV/c and LH2, C, Be, Bi, U targets.
- MIPP has collected 1.6×10^6 events of 120 GeV p striking the MINOS target.

Status of MIPP Analysis



- π^-/π^+ , K^-/K^+ , and K/π production ratios above 20 GeV/c agree well with expectations from MINOS beam-tuning.
- The MIPP Collaboration has completed the calibration of all PID detectors and is now focusing on the hadron production measurement from the NuMI target data set.
- See poster by Yusuf Gunaydin.



NuMI Beam



ν_μ CC Analysis


Precision measurement of
 Δm^2 and $\sin^2(2\theta)$

ν_μ CC Event Selection






- CC/NC separation achieved via a kNN event selection based on:
 - Track length
 - Mean pulse height
 - Fluctuation in pulse height
 - Transverse track profile
- Cut on separation parameter maximizes CC selection efficiency and minimizes NC background.
- Good agreement between data and MC above the CC/NC separation parameter cut.

Expected Far Detector Spectrum




Line Source
at ND

Point Source
at FD




- Near detector spectrum is extrapolated to the far detector.
- Use MC to provide energy smearing and acceptance corrections.

Systematic Uncertainties




- Systematic uncertainties estimated by fitting modified MC in place of data.
- ν_μ CC measurement is statistics limited.
- Dominant uncertainties are:
 - ND/FD relative normalization (Δm^2)
 - Overall hadronic energy calibration (Δm^2)
 - NC background ($\sin^2(2\theta)$)



- These three systematic effects are included in the final fit as nuisance parameters.


FD Energy Spectrum/Performing the Fit



- FD energy spectrum is only looked at after performing:
 - low-level data quality checks
 - procedural checks
- 848 events observed in the FD
- 1065 ± 60 expected with no oscillations
- We fit the energy distribution to the oscillation hypothesis:


$$P(\nu_\mu \rightarrow \nu_\mu) = 1 - \sin^2(2\theta) \sin^2(1.27 \Delta m^2 L/E)$$

Contours



- **Constrained fit:**
 - $\Delta m^2 = (2.43 \pm 0.13) \times 10^{-3} \text{ eV}^2$ (68% CL)
 - $\sin^2(2\theta) > 0.90$ (90% CL)
 - $\chi^2/\text{ndof} = 90/97$
- **Unconstrained fit:**
 - $\Delta m^2 = 2.33 \times 10^{-3} \text{ eV}^2$
 - $\sin^2(2\theta) = 1.07$
 - $\Delta\chi^2 = -0.6$

Alternative Hypotheses



Decay:

$$P_{\mu\mu} = (\sin^2\theta + \cos^2\theta \exp(-\alpha L/E))^2$$

$$\chi^2/\text{ndof} = 104/97$$

$$\Delta\chi^2 = 14$$

Disfavored at 3.7σ

Decoherence:

$$P_{\mu\mu} = 1 - \frac{1}{2} \sin^2(2\theta) (1 - \exp(-\mu^2 L/2E))$$

$$\chi^2/\text{ndof} = 123/97$$


$$\Delta\chi^2 = 33$$

Disfavored at 5.7σ

NC Analysis


The search for sterile neutrinos

NC Event Selection in the ND




- Since NC events probe active flavors, a depletion of NC events in the FD can only be explained by ν_s .
- We select reconstructed “shower-like” (short) events that fall within a fiducial volume.

Measured Near Detector Spectrum



NC event selection efficiency is 90%, purity is 60%.

3-Flavor Analysis Results



Data/MC Comparison for $\theta_{13} = 0$

Energy Range (GeV)	0 - 3	0 - 120
Data	100	291
MC	115.16 ± 7.67	292.63 ± 15.02
Significance (σ)	1.15	0.10

- For $E_{\text{vis}} < 3 \text{ GeV}$, $f_{\text{NC}} < 35\%$ at 90% CL.
- For $E_{\text{vis}} < 120 \text{ GeV}$, $f_{\text{NC}} < 17\%$ at 90% CL.



Other Finalized Analyses

- **“Sudden stratospheric warmings seen in MINOS deep underground muon data”**: High-energy cosmic muon rate is strongly correlated to temperature changes in the upper atmosphere. MINOS has shown that (under)ground-based high statistics cosmic muon measurements are a new tool to be used in tracking meteorological phenomena in the upper atmosphere.
- **“Testing Lorentz Invariance and CPT Conservation with MINOS Near Detector Neutrinos”**: search for a sidereal signal in the MINOS ND. Upper limits set on individual SME Lorentz and CPT violating terms.
- **“Observation of deficit in NuMI neutrino-induced rock and non-fiducial muons in MINOS far detector and measurement of neutrino oscillation parameters”**: see poster by Aaron McGowan

ν_e CC Analysis


The search for ν_e appearance

ν_e Background Estimates



- Measurement dominated by backgrounds: at the CHOOZ limit, 12 ν_e events are expected with 42 background events (for 3.25×10^{20} POT).
- Dominant background is NC events, with other significant contributions coming from ν_μ CC events and intrinsic beam ν_e events.
- We see a very large discrepancy between selected ν_e ND MC and data events.
- Two new data-driven methods have been developed to resolve the MC/data difference - see [posters by Steven Cavanaugh and Lisa Whitehead](#) for details.

ν_e Sensitivity



- Projected limits for expected MINOS integrated exposures for the next few years.
- Inverted hierarchy (in red) shown only for lowest exposure.
- MINOS can improve upon the CHOOZ limit by $\sim x2$.

Other Analyses in the Works

- Anti-neutrino oscillation measurements
- ND measurements:
 - Inclusive CC cross-section and structure functions
 - M_A extraction from quasi-elastic events
 - NC coherent scattering on Fe
 - Cosmic ray studies

Conclusions



- 2007-08 has been a very productive year for MINOS!
- Latest ν_μ CC analysis results (3.36×10^{20} POT):
 - $\Delta m^2 = (2.43 \pm 0.13) \times 10^{-3} \text{ eV}^2$ (68% CL),
 - $\sin^2(2\theta) > 0.90$ (90% CL),
 - Decay and decoherence models disfavored at 3.7 and 5.7 σ respectively.
- NC analysis results (2.46×10^{20} POT): fraction of disappearing NC events < 0.17 at 90% CL.
- Great progress in understanding the backgrounds and systematics in the ν_e appearance measurement; first results are expected later this year.
- Results from MIPP expected later this year, expected uncertainty on ν flux is $\sim 15\%$.
- Many ND ν interaction measurements also expected later this year.
- Thanks to FNAL AD, CD, and administration for all their hard work and support!

Backup Slides

Outline of the Rest of this Talk

- The MINOS Experiment
- What's new (2007-08)
- Future Prospects


MIPP Performance




- Momentum resolution is $\sim 5\%$ at 120 GeV/c, much better at lower momenta.
- Vertex resolution is ~ 8 mm in the beam direction, ~ 2 mm transverse.
- Reconstructed momentum appears to be systematically low by $\sim 2\%$.

MIPP Performance

ToF Velocity vs. Momentum, All Bars



TPC $\langle dE/dx \rangle$ vs. Momentum







- Ckov has ~ 5 pe per $\beta=1$ particle.
- ToF resolution is ~ 300 ps
- TPC $\langle dE/dx \rangle$ resolution is ~ 12 %.

ν_μ CC/NC Separation



- CC/NC separation achieved via a kNN

event selection based on:

- Track length
- Mean pulse height
- Fluctuation in pulse height
- Transverse track profile






ν_μ CC Event Selection






- Cut on separation parameter maximizes CC selection efficiency and minimizes NC background.
- Good agreement between data and MC above the CC/NC separation parameter cut.

Far Detector Low-level Data Quality Checks


- FD energy spectrum is only looked at after performing:
 - low-level data quality checks
 - procedural checks



ND Distributions After Making PID Cut




Sensitivity






- Final contour is a bit smaller than the predicted sensitivity because $\sin^2(2\theta)$ falls in the unphysical region.
- A study shows that 26.5% of unconstrained fits have a fit value of $\sin^2(2\theta) \geq 1.07$
- Feldman-Cousins study indicates that our contours are slightly conservative.

LE1 vs. LE2 Beam Configurations



NC Event Selection in the FD

- Identical cuts are made in FD as in ND.
- MC oscillated with 2007 MINOS CC best fit values of $\Delta m^2 = 2.38 \times 10^{-3} \text{ eV}^2$ and $\sin^2(2\theta) = 1$.




—•— Far Detector Data

— Monte Carlo



Cosmic Rays and Upper Atmospheric Weather

Sudden Stratospheric Warmings



- There is a strong correlation between the high energy cosmic ray rate and temperature changes in the upper atmosphere.
- The MINOS FD observes a large cosmic muons rate and can measure these percent-level changes in rate.
- SSWs have been tracked using balloon measurements, rocket soundings, LIDAR, airborne and satellite observations. MINOS now provides another new tool with which to observe these meteorological phenomena.

ν_e Data-Driven Background Studies



Estimate	Signal ν_e	Total BG	NC	ν_μ CC	Beam ν_e	ν_τ CC
Horn On/Off	12	42	29	8	3	2
MRCC	12	43	32	6	3	2

$\sin^2(2\theta_{23}) = 1.0$
 $\Delta m^2_{32} = 2.4 \times 10^{-3} \text{ eV}^2$
 $\sin^2(2\theta_{13}) = 0.15$
 no matter effects
 $3.25 \times 10^{20} \text{ POT}$

- Horn On/Off - constrain the relative ratios of NC and ν_μ CC background events in two different beam configurations.
- Muon removed hadron showers from ν_μ CC (MRCC).

QUESTIONS I HAVE

- Can someone remind me why the best fit value on the top (small) plot on slide 17 has $\sin^2(2t) > 1$? Is it just statistics? This plot was made with MC...
- Will the NC result be redone with our latest values of Δm^2 and $\sin^2(2\theta)$?
- Should I mention the Horn 1 problem?
- Can I mention the Nature article submission?

NOTE:

- I intend to add more backup slides, but if you have any specific suggestions, please let me know!